

The ILAE Classification of Seizures & the Epilepsies: Modification for Seizures in the Neonate. Proposal from the ILAE Task Force on Neonatal Seizures

Ronit M Pressler^{1,2}, Maria Roberta Cilio³, Eli M Mizrahi⁴, Solomon L Moshé⁵, Magda L Nunes⁶, Perrine Plouin⁷, Sampsa Vanhatalo⁸, Elissa Yozawitz⁹, Sameer M Zuberi¹⁰

- ¹ Clinical Neuroscience, UCL-Institute of Child Health, London, UK, r.pressler@ucl.ac.uk
² Department of Clinical Neurophysiology, Great Ormond Street Hospital for Children NHS Foundation Trust, London, UK
³ Departments of Neurology and Pediatrics, Benioff Children's Hospital, University of California San Francisco, San Francisco, CA, USA, roberta.cilio@ucsf.edu
⁴ Departments of Neurology, Baylor College of Medicine, Houston, TX, emizrahi@bcm.edu
⁵ Saul R. Korey Department of Neurology, Dominick P. Purpura Department of Neuroscience and Department of Pediatrics, Albert Einstein College of Medicine and Montefiore Medical Center, Bronx, New York, USA, Solomon.moshe@einstein.yu.edu
⁶ PUCRS School of Medicine, Porto Alegre, RS, Brazil, mlnunes@pucrs.br
⁷ Department of Clinical Neurophysiology, Hospital Necker Enfant Malades, Paris, France perrine.plouin@gmail.com
⁸ Department of Clinical Neurophysiology, Children's Hospital, HUS Imaging, Helsinki University Central Hospital and University of Helsinki, Finland, sampsa.vanhatalo@helsinki.fi
⁹ Saul R. Korey Department of Neurology and Department of Pediatrics, Albert Einstein College of Medicine and Montefiore Medical Center, Bronx, New York, USA eyozawit@montefiore.org
¹⁰ Paediatric Neurosciences Research Group, Fraser of Allander Neurosciences Unit Royal Hospital for Sick Children, Glasgow, UK, sameer.zuberi@nhs.net

Corresponding author:

Dr Ronit M Pressler
Department of Clinical Neurophysiology
Great Ormond Street Hospital for Children NHS Foundation Trust
London WC1N 3JH, UK
Tel +442078138471
ronit.pressler@gosh.nhs.uk

Keywords: Neonatal seizures, EEG, semiology, classification, epilepsy

Text pages: 16 (incl. abstract, definitions, and disclosures, excl. references)
References: 99
Figures: 3
Tables: 5

Abstract

Seizures are the most common neurological emergency in the neonatal period occurring in 1–5 per 1000 live births. In contrast to seizures in infancy and childhood, most seizures in the neonate are symptomatic of an acute illness and can be electrographic only or associated with clinical manifestations. Hence, neonatal seizures may not fit easily into classification schemes for seizures and epilepsies primarily developed for older children and adults. The Task Force on Neonatal Seizures was established with the aim of developing a modification of the 2017 ILAE Classification of Seizures and Epilepsies, relevant to neonates. The proposed neonatal classification framework emphasizes the role of EEG in the diagnosis of seizures in the neonate and includes a classification of seizure types relevant to this age group. The seizure type is typically determined by the predominant clinical feature. As seizures in neonates are often electrographic only with no clinical feature, these are included in the proposed classification. Clinical events without an EEG correlate are not included. As seizures in the neonatal period have been shown to have a focal onset, a division into focal and generalized is unnecessary. Seizures can have a motor (automatisms, clonic, epileptic spasms, myoclonic, sequential, tonic) or non-motor (autonomic, behavior arrest) presentation. The proposed classification allows the user to choose the degree of detail when classifying seizures in this age group.

The proposed classification was successfully applied to 147 video-EEG recordings of neonatal seizures. These included representative samples of the common etiologies; hypoxic ischemic encephalopathy, stroke, infection, metabolic and genetic disorders. We noted that certain seizure types were associated with particular etiologies emphasizing the value in describing seizure types in this age group.

Keywords: Neonatal seizures, EEG, semiology, classification, epilepsy

Key points:

- A new framework is proposed for seizures in neonates, in keeping with 2017 ILAE seizure classification while tailored to the neonatal period
- The framework emphasizes the necessity of EEG diagnosis of seizures in the neonatal period.
- Seizures can occur with clinical manifestations (motor or non-motor) or without clinical manifestations (electrographic only).
- Descriptors are determined by the predominant clinical feature and divided into motor and non-motor.

Definitions

For the purpose of this report, the following definitions are used:^{1,2}

- Gestational age (GA): time elapsed between the first day of the last menstrual period and the day of deliver (completed weeks).
- Postmenstrual age (PMA): gestational age plus chronological age (in weeks).
- Conceptional age (CA): age from conception: PMA minus 2 weeks.
- Preterm infant: born before GA of 37 weeks.
- Neonatal period: Period from birth up to and under 28 days in term neonates, or from birth up to a PMA of 40 weeks and under 28 days in preterm neonates.

Introduction

The 2017 ILAE Position Papers on Classification of Seizure Types and the Epilepsies^{3,4} presented a framework for classification including seizure types, epilepsy types, and syndromes. There is an emphasis on defining etiology at all levels of clinical classification in addition to consideration of comorbidities. Terminology is updated with some new seizure types included. The ILAE Commission on Classification & Terminology recognized that seizures in the neonate require special considerations and therefore a Neonatal Task Force was established with the aim of integrating seizures and epilepsies in this age group into the 2017 ILAE Classification.

Seizures are the most common neurological emergency in the neonatal period occurring in 1–5 per 1000 live births. The majority of neonatal seizures are symptomatic of an acute illness with an underlying etiology either documented or suspected.^{5,6} Epilepsy syndromes may present in the neonatal period and with the increasing availability of genetic testing expanding numbers of neonatal epilepsies with genetic and metabolic etiologies are recognized.⁷ Although many causes can give rise to neonatal seizures, a relatively small number account for most seizures (Figure 1).

The clinical diagnosis of neonatal seizures is difficult, particularly in critically ill infants due to the multitude of epileptic and non-epileptic clinical manifestations within the intensive care setting.^{12,13} In the study by Malone,¹⁴ 20 video clips of paroxysmal events in neonates were presented to 137 health professionals (mostly neonatologists and intensivists) with the aim of classifying movements as seizure or non-seizure. Only 50% of events were correctly classified. There was poor inter-observer agreement independent of observers' specialty. The immature state of the motor pathways^{15,16} in term and preterm neonates may account for some of the difficulty in differentiating seizures from non-epileptic movements.¹⁷ In

1
2
3 selected populations, particularly in infants with hypoxic-ischemic encephalopathy (HIE), 50-
4
5 80% of seizures are electrographic only and, as a result, the extent of the seizure burden
6
7 (time spent with seizures; defined as electrographic seizures in minutes per hour) may be
8
9 greatly underestimated.^{6,12-14,18} It has been suggested that hypothermia therapy for HIE
10
11 may also increase electro-clinical uncoupling of seizures, with a high incidence of
12
13 electrographic seizures in these infants.¹² There is evidence that electrographic seizure
14
15 burden has a comparable effect on outcome as electro-clinical seizures.¹⁹⁻²³

16
17
18 The ILAE defines a seizure as a transient occurrence of signs and/or symptoms due to
19
20 abnormal excessive or synchronous neuronal activity in the brain;^{3,24} however, a seizure
21
22 does not necessarily mean that a person has epilepsy. Electrographic only seizures are not
23
24 included in this definition.

25
26 Neonatal seizures are often categorized as clinical only, electro-clinical, or electrographic-
27
28 only.^{13,25} A clinical only seizure consists of a sudden paroxysm of abnormal clinical changes
29
30 without a definite EEG association. Whether or not all of these events are indeed of
31
32 epileptic origin remains uncertain in newborns. An electro-clinical seizure features definite
33
34 clinical signs simultaneously coupled with an electrographic seizure. An electrographic only
35
36 seizure refers to the presence of a definite EEG seizure that is not associated with any
37
38 evident clinical signs (synonyms: clinically silent or subclinical seizures). The term
39
40 electrographic only is preferred as this depends on observational methods used and the
41
42 seizure may not be truly subclinical.

43
44
45 The American Clinical Neurophysiology Society has recently defined an electrographic
46
47 neonatal seizures as a paroxysmal abnormal, sustained change in the EEG, characterized by
48
49 a repetitive and evolving pattern with a minimum 2 μ V voltage (peak to peak) and duration
50
51 of at least 10 seconds.²⁶ This definition does not require any evident clinical change.
52
53
54
55
56
57
58
59
60

1
2
3 However, this definition has been challenged as brief runs of rhythmic discharges (BRD) or
4
5 brief rhythmic ictal discharges (BIRDs) were shown to be associated, like electrographic
6
7 seizures, with clinical manifestations and poor outcome.^{27,28}
8
9

10 11 12 13 14 15 **Historical review**

16
17
18 Historical efforts to characterize and classify neonatal seizures have been directed towards
19
20 emphasizing how they differ from those of older children and adults. In this report our aim
21
22 is to use terminology consistent with the 2017 ILAE Classification of Seizures and the
23
24 Epilepsies.^{3,4,24}

25
26
27 Studies in the 1950's and early 1960's focused on motor and behavioral changes and were
28
29 based upon direct observation with or without EEG recordings.²⁹⁻³¹ This resulted in the use
30
31 of terms such as focal clonic and generalized tonic for neonatal seizure classification
32
33 (generalized tonic-clonic seizures were considered non-existent in this age group).
34
35

36
37 Eventually, myoclonus was added to the growing classification.³²

38
39
40 Early investigators recognized autonomic nervous system changes including variation in
41
42 respiratory rate, vasomotor changes, salivation, heart rate and blood pressure as seizure
43
44 manifestations. Polymorphic and atypical clinical events were described, the latter including
45
46 staring, sudden awakening and alerting, eye deviation, eye blinking, nystagmus, chewing
47
48 and limb movements resembling "swimming, rowing, and pedaling".³³ These descriptions
49
50 were incorporated into classifications schemes as a category of "anarchic",³¹ "minimal"³⁴ or
51
52 "subtle"³⁵. These findings resulted in the classification proposed by Volpe which included:
53
54 multifocal clonic, focal clonic, tonic, myoclonic and subtle seizures.^{35,36}
55
56
57
58
59
60

1
2
3 Using EEG-video analysis of electro-clinical correlations,³⁷ it was recognized that some
4
5 clinical events previously reported as seizures were in fact non-epileptic. Indeed, many of
6
7 these clinical events were found to occur without EEG correlate and could be provoked by
8
9 stimulation and suppressed by restraint.¹³ This led to a reconsideration of the classification
10
11 of neonatal seizures based upon either pathophysiology (epileptic versus non-epileptic);
12
13 electro-clinical relationships (electro-clinical, clinical only, electrical only); or behavioral
14
15 (focal clonic, focal tonic, myoclonic, spasms, generalized tonic, motor automatisms – each
16
17 with additional modifiers to suggest whether they were considered to be of epileptic or
18
19 non-epileptic origin). The term motor automatisms included: ocular movements, oral-
20
21 buccal-lingual movement, and progression movements of the limbs (pedaling, swimming,
22
23 rowing). The basic movements and behaviors of earlier classification systems were
24
25 preserved and epileptic spasms were added. In addition, a significant group of neonatal
26
27 seizures were considered to have only an electrographic signature with no clinical
28
29 correlate.¹³

30
31
32 With the advent of prolonged bedside electrographic monitoring in the Neonatal Intensive
33
34 Care Unit (NICU), either with EEG or with amplitude integrated EEG (aEEG), it has been
35
36 increasingly recognized that electrographic seizures without clinical correlates are frequent,
37
38 particularly in critically-ill neonates.^{18,38,39} As a result, the definition of neonatal seizures has
39
40 been reconsidered, now with a focus on the electrographic basis of the events, either with
41
42 or without clinical manifestations.⁴⁰

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Proposal

The goal of this report is to propose a classification of seizures in neonates that can fulfil the following criteria:

- Integrate into the 2017 ILAE Classification
- Be based on electro-clinical phenotype
- Emphasize the key role of EEG in the diagnosis of neonatal seizures
- Have implication for management and treatment of events
- Be acceptable to neonatologists, pediatricians, epileptologists, neurophysiologists, and neurologists.
- Be applicable in all health care settings.

Figure 2 depicts the proposed approach for seizure classification.

Presentation

Newborns may present with paroxysmal clinical events suspected to be epileptic seizures; these include motor or non-motor phenomena.

Diagnosis

In neonates, video-EEG recording is the gold standard for diagnosis.^{13,41-43} A proportion of seizures are electrographic only, particularly in encephalopathic and critically ill patients.

^{12,18,38,44} In the neonate, this may occur because of the immaturity of the central nervous system. Uncoupling may increase after administration of antiseizure medications, particularly phenobarbital.^{13,18,39,45-47} Therefore, electrographic-only seizures should to be part of the classification. The initial stage of description of a neonatal seizure should specify

whether a seizure is with (electro-clinical) or without clinical signs (electrographic only).

Electro-clinical dissociation is the phenomenon when a clinical seizure type occurs at times with and at times without an associated rhythmic EEG discharge in a given patient.^{45,46}

However this is considered to be a rare occurrence and by definition implies that electrographic seizures (with or without clinical correlate) also occur in that given patient.

Therefore, only events with EEG correlate are included in this proposal.

Seizure types

We used the definition of seizure type as suggested by Fisher and colleague,³ a useful grouping of seizure characteristics for purposes of communication in clinical care, teaching, and research.

The basic principles of the 2017 ILAE classification of seizure types³ (see appendix Figure A) are based on the 1981 classification with the initial division of seizures into those of focal and generalized onset.^{48,49} As newborns have been shown to have seizures with exclusively focal onset,^{13,50} the initial division into focal and generalized is unnecessary. The second step in the 2017 ILAE classification is the division into aware and unaware (impaired awareness) seizures, however, this is not applicable to neonates as it is not possible to confidently and reproducibly assess awareness and responsiveness in this age group.

This is followed by the division into motor and non-motor seizures and finally by the seizure type (Table 1). In the 2017 ILAE classification, focal seizures are determined by the first feature, as seizure localization has important clinical implications to identify a possible focal lesion irrespective of etiology. While seizures in neonates can present with a variety of clinical signs, in the majority of cases a single predominant feature can be determined.

Pragmatically, it appears best to classify seizures according to the predominant clinical

1
2
3 manifestation, as this is more likely to have clinical implications for etiology than
4
5
6 determination of the seizure onset zone. This may or may not be the first clinical
7
8 manifestation. For example, a neonate may present with a focal tonic posturing and in
9
10 addition have some ocular myoclonus – this can still be classified as a tonic seizure.
11
12 In some situations, it may be difficult to identify the dominant feature, typically in longer
13
14 seizures where a sequence of clinical features can be seen, often with changing
15
16 lateralization. Events with a sequence of signs, symptoms, and EEG changes at different
17
18 times have been described as a sequential seizure in the 2017 ILAE classification manual⁵¹.
19
20 As this is often seen in neonates this term was added to the seizure types.
21
22 Several seizure types described in the 2017 ILAE classification^{3,51} cannot be diagnosed in
23
24 newborns due to lack of verbal and limited non-verbal communications. These include
25
26 sensory seizures, cognitive and emotional seizures. Sensory seizures are defined as a
27
28 perceptual experience not caused by appropriate stimuli in the external world. Such seizures
29
30 can in rare cases produce reproducible semiology but it is assumed that in the majority of
31
32 cases they would appear as electrographic only events. Awareness and responsiveness
33
34 cannot be accurately assessed in neonates and hence not readily classified; however, this
35
36 may change with more advanced technology or detailed observation. Similarly,
37
38 somatosensory or visual auras cannot be determined in neonates. Due to the relative low
39
40 muscle tone and supine position of newborns, the occurrence of atonic seizures cannot be
41
42 evaluated clinically without invasive methods. These seizure types are therefore not
43
44 included in the proposed framework. Motor seizures can be further described using
45
46 modifiers listed in Table 2. The framework allows the user to classify the seizure in as much
47
48 detail as required in a certain situation. The full description would include manifestation, a
49
50
51
52
53
54
55
56
57
58
59
60 descriptor and etiological diagnosis.

Epilepsy syndromes

While the majority of seizures in the neonatal period occur in the context of an acute illness, in some cases the seizures may be the first manifestation of early-onset epilepsy. Early differentiation of acute symptomatic seizures from neonatal-onset epilepsies has important therapeutic and prognostic implications since the evaluation and long-term management of neonatal epilepsies are distinct from those of acute symptomatic seizures.⁷ Syndromes presenting in the neonatal period include: self-limited (benign) familial neonatal epilepsy, early myoclonic encephalopathy (EME), and early infantile epileptic encephalopathy (Ohtahara syndrome).⁷⁰

Recent advances in neuroimaging and genomic technology as well as the implementation of video-EEG in the NICU, allow for the identification of more discrete, etiology-specific neonatal epilepsy syndromes than previously recognized.⁷¹ It is likely that the combination of more sophisticated genetic testing and video-EEG monitoring will allow the identification and stratification of distinct etiology-specific electro-clinical phenotypes,⁷² as suggested in the new ILAE classification of the epilepsies (Figure 3).⁴

Evaluation of the proposal

Methods

During a 2-day workshop and face-to-face meetings the classification and framework were tested by reviewing of video-EEG data of neonatal seizures. Inclusion criteria for the data collection were as follows: (1) Consecutive and unselected cases from neonatal units at Albert Einstein College of Medicine, Baylor College of Medicine, Helsinki University Central Hospital, PUCRS School of Medicine, Royal Hospital for Children Glasgow, UCL-Institute of

1
2
3 Child Health, and University of California San Francisco, and (2) Adequate EEG and video
4
5
6 quality to verify ictal event and to review semiology. In order to represent common
7
8 etiologies, we decided to include a minimum of 30 cases with HIE, 20 with stroke or
9
10 hemorrhages, 20 with infectious causes, 20 with inborn errors of metabolism, 20 with
11
12 genetic causes, and 10 with cortical malformations.

13
14
15 A selection bias is recognized as most of the centers are tertiary referral centers with or
16
17 without attached maternity hospitals. The proportion of genetic, structural and metabolic
18
19 etiologies may be somewhat higher than seen in other neonatal units.

20
21
22 The videos of all cases were reviewed by the members of the task force, the semiology
23
24 evaluated and the seizure classified based on the proposed classification.

25
26
27 An electrographic seizure was defined as an electrographic event with a pattern
28
29 characterized by sudden, repetitive, evolving stereotyped waveforms of with a beginning
30
31 and end which may or may not be accompanied by paroxysmal clinical changes. No
32
33 minimum duration was specified as long as there was sufficient demonstration of evolution
34
35 in frequency and morphology of the discharge. The exceptions to the concept of evolving
36
37 waveforms are clinical seizures such as myoclonic seizures and spasms which are associated
38
39 with an EEG correlate that is very brief and not evolving. The EEG was used to confirm that
40
41 an event was an epileptic seizure but not further analyzed.
42
43
44
45
46
47
48
49

50 **Results**

51
52 A total of 157 events from 146 neonates were reviewed. Ten infants were excluded because
53
54 of poor video quality (n=7) or EEG did not conclusively confirm a seizure (n=3), leaving a
55
56 total of 147 seizures in 136 neonates. Demographics are summarized in Table 3. The most
57
58
59
60

1
2
3 common etiology was HIE in term infants (25%) and vascular (stroke or hemorrhage) in
4
5 preterm infants (45%).
6
7

8 We were able to classify all seizures according to the proposed classification. Table 4 details
9
10 the seizure type according to etiology. The most common seizure type was electrographic
11
12 only both in term (45/131 seizures) and preterm infants (14/20 seizures). The most common
13
14 etiologies of electrographic seizures were HIE and infection. Clonic seizures were typically
15
16 seen in association with vascular etiologies (term infants only). Tonic seizures and sequential
17
18 seizures were most commonly observed in genetic etiologies and myoclonic seizures in
19
20 inborn errors of metabolism. All seizures in neonates with a genetic etiology had motor
21
22 manifestations. Automatism and behavioral arrest were uncommon as dominant seizure
23
24 manifestation. In preterm infants 70% (14/20) of seizures were electrographic regardless of
25
26 etiology. No dominant seizure type was seen for acute metabolic disorders, cortical
27
28 malformations and unknown causes but numbers were too small for comparison.
29
30
31
32
33
34
35
36
37
38
39

40 Discussion

41
42 Seizure semiology is the description of signs and symptoms associated with an ictal event
43
44 and is valuable in localizing the epileptogenic zone. However, in the neonate, the
45
46 development within the limbic system with its connections to midbrain and brainstem is
47
48 more advanced than the cerebral cortical organization,⁷³ leading to a higher frequency of
49
50 oral automatisms, ocular changes such as eye deviation, apnea and clinical features related
51
52 to the autonomic nervous system in neonates than in older children.
53
54

55
56 The neonatal classification proposed emphasizes the role of EEG in the diagnosis of seizures
57
58 in the neonate and includes a classification of seizure types relevant to this age group. The
59
60

1
2
3 seizure type is typically determined by the predominant clinical feature while in the 2017
4
5 ILAE classification of seizures in other age groups, the seizure type was defined by the first
6
7 manifestation. This was based on the assumption that the first manifestation may indicate
8
9 whether the seizure may have a focal onset. Since all seizures in the neonate are focal, we
10
11 propose to emphasize the predominant feature as this may provide clues regarding the
12
13 etiology.
14
15
16

17
18 This concept was evaluated by the Task Force by reviewing 147 neonatal seizures. We
19
20 demonstrated that the classification can be implemented in clinical practice. Among the
21
22 seizures reviewed we noted that certain etiologies are associated with certain seizure types:
23
24 most neonates with genetic etiologies have tonic seizures and / or sequential seizure,
25
26 whereas in preterm infants (regardless of etiology) and neonates with HIE or infectious
27
28 causes present most commonly with electrographic only seizures. Term infants with stroke
29
30 are more likely to have focal clonic seizures. Myoclonic seizures are associated with inborn
31
32 errors of metabolism.
33
34
35

36
37 Increased movements were noticed at arousal but we were not able to differentiate these
38
39 from other arousals either on clinical grounds or by EEG. However, hyperkinetic phases
40
41 were observed as part of sequential seizures. It was concluded that hyperkinetic events if
42
43 occurring in isolation cannot be defined as seizures, but they may occur in the context of a
44
45 sequential seizures.
46
47
48

49
50 Review of the literature and our own data suggest that seizure semiology in neonates may
51
52 have diagnostic value in respect to etiology of seizures and /or outcome (see Table 5).
53

54
55 However, many of these clinical associations are based on small case studies or with very
56
57 limited description of semiology and will need to be tested on a larger dataset.
58
59
60

1
2
3 Clancy and Legido described electrographic only seizures in newborns as sudden, repetitive,
4
5 evolving stereotyped waveforms with a definite beginning, middle, and end and a minimum
6
7 duration of 10 seconds.³⁸ However, the choice of 10 seconds duration was explicitly
8
9 arbitrary. Similarly an arbitrary minimum duration of 10 seconds is also applied to the
10
11 definition of a seizure in critical ill adults.⁸⁷ This is in contrast to some electro-clinical
12
13 seizures such as myoclonic seizures or spasms which are by definition shorter than 10
14
15 seconds.^{3,51,88} Both in neonates and critically ill adults it has been suggested that rhythmic
16
17 discharges of less than 10 seconds duration (so called BIRDs: Brief Interictal Rhythmic
18
19 Discharges) are associated with seizures in the same or subsequent EEG recording^{27,28,89,90}
20
21 and an increased risk of abnormal neurodevelopmental outcome.²⁷ BIRDs are defined as
22
23 very brief (<10 seconds) runs of focal or generalized sharply contoured rhythmic activity,
24
25 with or without evolution, that are not consistent with any known normal or benign pattern,
26
27 which in adults have a frequency greater than 4 Hz.⁹¹ It is of interest that in these clinical
28
29 studies BIRDs either showed no evolution^{90,91} or the presence or absence of evolution has
30
31 not been described.^{27,28} It has been suggested that definite BIRDs with an evolution
32
33 represent “very brief” electrographic seizures.⁸¹

34
35 We propose to define seizures in the neonatal period as an electrographic event with a
36
37 pattern characterized by sudden, repetitive, evolving stereotyped waveforms with a
38
39 beginning and end. However the duration has to be sufficient to demonstrate evolution in
40
41 frequency and morphology of the discharges which will depend on the frequency of the
42
43 discharge. The duration of this is to an extent arbitrary but need to be sufficient to allow
44
45 recognition of onset, evolution and resolution of an abnormal discharge. **BIRDs without**
46
47 **evolution are not considered seizures but may serve as an early predictor of seizures during**
48
49 **subsequent EEG monitoring and as a prognostic indicator.** Notable exceptions are certain
50
51
52
53
54
55
56
57
58
59
60

1
2
3 clinical seizures such as myoclonic seizures and spasms which are associated with an EEG
4
5 correlate that is very brief and not evolving. We acknowledge that with advanced
6
7 technology this definition may change in the future.
8
9

10 In defining clinical and electrographic seizures, we acknowledge that treatment may alter
11
12 both semiology and EEG features of these events, so that our definitions apply to seizure
13
14 recorded prior antiepileptic drug therapy. Electrographic seizure burden and seizure
15
16 frequency may impact the treatment approach, but the presence or absence of clinical signs
17
18 should not.
19
20
21

22
23 It is recognized that clinical care for affected neonates is also provided in centers where
24
25 resources and expertise are limited.⁹² EEG is considered the gold standard for seizure
26
27 diagnosis in the neonate, but in situations when and where it is not readily available, aEEG
28
29 may be used,⁹³ although its limitations are well recognized.^{94,95} If neither is available, we
30
31 would like to refer to the algorithm for diagnosis and treatment of neonatal seizures in
32
33 developing countries⁹² which was developed as part of the Global Campaign against
34
35 Epilepsy, an initiative by the ILAE and the World Health Organization (WHO). This document
36
37 proposes pragmatic guidelines for the diagnosis and management of the most common and
38
39 important conditions that may cause seizures in the neonatal period in centers with limited
40
41 resources, where recommended diagnostic testing may not be possible. In such situations,
42
43 clonic seizures may be identified more reliably than other seizure types¹⁴ as well as focal
44
45 tonic seizures and some sequential seizures whereas automatisms, autonomic seizures and
46
47 seizures with behavioral arrest require EEG confirmation and electrographic seizures will, by
48
49 definition, be missed without EEG. Beside maneuvers such as stimulation of the infant to
50
51 provoke behaviors similar to spontaneously observed clinical event suspected of being
52
53 seizures and restraint of infant limbs during spontaneous events to arrest the events can
54
55
56
57
58
59
60

1
2
3 help in identifying clinical events as exaggerated reflex behaviors and non-epileptic in
4
5 origin.¹³
6
7

8 Although this framework was developed for seizures in the neonatal period, we believe that
9
10 it can be readily applied to acute symptomatic seizures in critically ill patients of any age,
11
12 particularly within the intensive care setting who may present with similar manifestations.
13
14 Non convulsive seizures are common in critically ill patients⁹⁶ and electrographic only
15
16 presentation due to electro-clinical uncoupling has been described in 2/3 of critically ill
17
18 children with seizures.⁹⁷⁻⁹⁹ However, the etiologies may vary with age.
19
20
21
22
23
24
25
26
27

28 **Acknowledgements**

29
30 The workshop was partially funded by the ILAE. We would like to thank Dr Marie-Coralie
31
32 Cornet, MD (San Francisco, US) for her help in selecting and collecting video-EEGs used for
33
34 this study.
35
36
37
38
39
40
41
42
43

44 **Disclosure**

45
46
47 Ronit Pressler has no conflicts of interest in regards to this article. She receives consultant
48
49 fees from UCB.
50
51

52 Solomon L. Moshé is the Charles Frost Chair in Neurosurgery and Neurology and partially
53
54 funded by grants from NIH U54 NS100064 and NS43209, US Department of Defense
55
56 (W81XWH-13-1-0180 and EP170020), CURE Infantile Spasms Initiative and the Heffer Family
57
58 and the Segal Family Foundations and the Abbe Goldstein/Joshua Lurie and Laurie Marsh/
59
60

1
2
3 Dan Levitz families. He has no conflicts of interest in regards to this article. He is serving as
4
5 Associate Editor of Neurobiology of Disease and is on the editorial board of Brain and
6
7 Development, Pediatric Neurology and Physiological Research. He receives from Elsevier an
8
9 annual compensation for his work as Associate Editor in Neurobiology of Disease and
10
11 royalties from 2 books he co-edited. He received a consultant fee from Eisai, Mallinkrodt
12
13 and UCB.
14
15
16

17
18 Eli M. Mizrahi has no conflicts of interest in regards to this article. He has received
19
20 consultant fees from Eisai, Inc and royalties from Elsevier, McGraw-Hill and Springer
21
22 publishers.
23
24
25

26 The other authors have no conflict of interest to disclose in relation to this publication. We
27
28 confirm that we have read the Journal's position on issues involved in ethical publication
29
30 and affirm that this report is consistent with those guidelines.
31
32
33

34 35 36 37 **Ethical Publication Statement** 38

39
40 We confirm that we have read the Journal's position on issues involved in ethical publication
41
42 and affirm that this report is consistent with those guidelines.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Engle WA. American Academy of Pediatrics Committee on Fetus and Newborn. Age terminology during the perinatal period. *Pediatrics* 2004;114:1362–1364.
2. WHO. Preterm birth, fact sheet. 2016.
<http://www.who.int/mediacentre/factsheets/fs363/en/>.
3. Fisher RS, Cross JH, French JA, et al. Operational classification of seizure types by the International League Against Epilepsy: Position Paper of the ILAE Commission for Classification and Terminology. *Epilepsia* 2017;58:522-530.
4. Scheffer IE, Berkovic S, Capovilla G, et al. ILAE classification of the epilepsies: Position paper of the ILAE Commission for Classification and Terminology. *Epilepsia* 2017;58:512-521.
5. Ronen G, Penney S, Andrews W. The epidemiology of clinical neonatal seizures in Newfoundland: A population-based study. *J Pediatr* 1999;134:71-75.
6. Glass H, Shellhaas RA, Wusthoff CJ, et al. Contemporary Profile of Seizures in Neonates: A Prospective Cohort Study. *J Pediatr* 2016;174:98-103.
7. Shellhaas RA, Wusthoff CJ, Tsuchida TN et al. Profile of neonatal epilepsies: Characteristics of a prospective US cohort. *Neurology* 2017;89:893-899.
8. Levene MI, Trounce JQ. Cause of neonatal convulsions. Towards more precise diagnosis. *Arch Dis Child* 1986;61:78–79.
9. Lanska MJ, Lanska DJ, Baumann RJ, et al. A population-based study of neonatal seizures in Fayette County, Kentucky. *Neurology* 1995;45:724-732.

- 1
2
3 10. Tekgul H, Gauvreau K, Soul J, et al. The current etiologic profile and neurodevelopmental
4
5 outcome of seizures in term newborn infants. *Pediatrics* 2006;117:1270-1280.
6
7
- 8
9 11. Janackova S, Boyd S, Yozawitz E, et al. Electroencephalographic characteristics of
10
11 epileptic seizures in preterm neonates. *Clin Neurophysiol* 2016;127:2721-2727.
12
13
- 14 12. Nash KB, Bonifacio SL, Glass HC, et al. Video-EEG monitoring in newborns with hypoxic-
15
16 ischemic encephalopathy treated with hypothermia. *Neurology* 2011;76:556-562.
17
18
- 19 13. Mizrahi EM, Kellaway P. Characterization and classification of neonatal seizures.
20
21 *Neurology* 1987;37:1837-1844.
22
23
- 24 14. Malone A, Ryan CA, Fitzgerald A, et al. Interobserver agreement in neonatal seizure
25
26 identification. *Epilepsia* 2009;50:2097-2101.
27
28
- 29 15. Haut SR, Velísková J, Moshé SL. Susceptibility of immature and adult brains to seizure
30
31 effects. *Lancet Neurol* 2004;3:608-617.
32
33
- 34 16. Galanopoulou AS, Moshé SL. In search of epilepsy biomarkers in the immature brain:
35
36 goals, challenges and strategies. *Biomark Med* 2011;5:615-628.
37
38
- 39 17. Murray DM, Boylan GB, Ali I, et al. Defining the gap between electrographic seizure
40
41 burden, clinical expression and staff recognition of neonatal seizures. *Arch Dis Child*
42
43 *Fetal Neonatal Ed* 2008;93:F187-F191.
44
45
- 46 18. Scher MS, Alvin J, Gaus L, et al. Uncoupling of EEG-clinical neonatal seizures after
47
48 antiepileptic drug use. *Pediatr Neurol* 2003;28:277-280.
49
50
- 51 19. McBride MC, Laroia N, Guillet R. Electrographic seizures in neonates correlate with poor
52
53 neurodevelopmental outcome. *Neurology* 2000;55:506-513.
54
55
56
57
58
59
60

- 1
2
3 20. Miller SP, Weiss J, Barnwell A, et al. Seizure-associated brain injury in term newborns
4
5 with perinatal asphyxia. *Neurology* 2002;58:542-548.
6
7
8
9 21. Boylan GB, Pressler RM, Rennie JM, et al. Outcome of electroclinical, electrographic and
10
11 clinical seizures in the newborn infant. *Dev Med Child Neurol* 1999;41:819–825.
12
13
14 22. Srinivasakumar P, Zempel J, Trivedi S, et al. Treating EEG Seizures in Hypoxic Ischemic
15
16 Encephalopathy: A Randomized Controlled Trial. *Pediatrics* 2015;136:e1302-1309.
17
18
19
20 23. Kharoshankaya L, Stevenson NJ, Livingstone V, et al. Seizure burden and
21
22 neurodevelopmental outcome in neonates with hypoxic-ischemic encephalopathy. *Dev*
23
24 *Med Child Neurol* 2016;58:1242-1248.
25
26
27
28 24. Fisher RS, van Emde Boas W, Blume W, et al. Epileptic seizures and epilepsy: definitions
29
30 proposed by the International League Against Epilepsy (ILAE) and the International
31
32 Bureau for Epilepsy (IBE). *Epilepsia* 2005;46:470-472.
33
34
35
36 25. Scher MS. Controversies regarding neonatal seizure recognition. *Epileptic Disord*
37
38 2002;4:139-158.
39
40
41
42 26. Tsuchida TN, Wusthoff CJ, Shellhaas RA, et al. American clinical neurophysiology society
43
44 standardized EEG terminology and categorization for the description of continuous EEG
45
46 monitoring in neonates: report of the American Clinical Neurophysiology Society critical
47
48 care monitoring committee. *J Clin Neurophysiol* 2013;30:161-173.
49
50
51
52 27. Oliveira AJ, Nunes ML, Haertel LM, et al. Duration of rhythmic EEG patterns in neonates:
53
54 new evidence for clinical and prognostic significance of brief rhythmic discharges. *Clin*
55
56 *Neurophysiol* 2000;111:1646-1653.
57
58
59
60

- 1
2
3 28. Nagarajan L, Palumbo L, Ghosh S. Brief electroencephalography rhythmic discharges
4
5 (BERDs) in the neonate with seizures: their significance and prognostic implications. J
6
7 Child Neurol 2011;26:1529-1533.
8
9
10
11 29. Burke JB. The prognostic significance of neonatal convulsions. Arch Dis Child
12
13 1954;29:342-345.
14
15
16
17 30. Harris R, Tizard JP. The electroencephalogram in neonatal convulsions. J Pediatr
18
19 1960;57:501-520.
20
21
22 31. Dreyfus-Brisac C, Monod N. Electroclinical studies of status epilepticus and convulsions
23
24 in the newborn: In Kellaway P, Petersen I (Eds) Neurological and
25
26 Electroencephalographic Correlative Studies in Infancy. New York: Grune and Statton,
27
28 1964:250-272.
29
30
31
32 32. Rose AL, Lombroso CT. A study of clinical, pathological, and electroencephalographic
33
34 features in 137 full-term babies with a long-term follow-up. Pediatrics 1970;45:404-425.
35
36
37
38 33. Minkowski A, Ste Anne-Dargassies S, Dreyfus-Brisac C, et al. [Convulsive state in the
39
40 newborn infant]. Arch Fr Pediatr 1955;12:271-284.
41
42
43
44 34. Lombroso CT. Seizures in the newborn. In Magnus O, Lorentz de AM (Eds) Handbook of
45
46 Clinical Neurology, The Epilepsies. Amsterdam: North Holland, 1974;15:189-218.
47
48
49
50 35. Volpe J. Neonatal seizures. New Engl J Med 1973;289:413-416.
51
52
53
54 36. Volpe JJ. Neonatal seizures: current concepts and revised classification. Pediatrics
55
56 1989;84:422-428.
57
58
59
60

- 1
2
3 37. Watanabe K, Hara K, Miyazaki S, et al. Electroclinical studies of seizures in the newborn.
4
5 Folia Psychiatr Neurol Jpn 1977;31:381-392.
6
7
8
9 38. Clancy RR, Legido A, Lewis D. Occult neonatal seizures. *Epilepsia* 1988;29:256-261.
10
11
12 39. Boylan GB, Rennie JM, Pressler RM, et al. Phenobarbitone, neonatal seizures and video-
13
14 EEG. *Arch Dis Child* 2002;86:F165-170.
15
16
17
18 40. Mizrahi EM, Pressler RM. Foundations of neonatal epileptology: classification of seizures
19
20 and epilepsies in the neonate and their aetiology, electroencephalography, prognosis
21
22 and pathophysiology. In Moshé SL, Cross JH de Bellescize J, et al (Eds) *Seizures of Onset*
23
24 *in the First Two Years of Life*. New York: McGraw Hill, 2015:8-27.
25
26
27
28 41. Shellhaas RA, Chang T, Tsuchida T, et al. The American Clinical Neurophysiology Society's
29
30 guideline on continuous electroencephalography monitoring in neonates. *J Clin*
31
32 *Neurophysiol* 2011;28:611-617.
33
34
35
36 42. Plouin P, Kaminska A. Neonatal seizures. *Handb Clin Neurol* 2013;111:467-476.
37
38
39
40 43. Silverstein FS, Jensen FE, Inder T, et al. Improving the treatment of neonatal seizures:
41
42 National Institute of Neurological Disorders and Stroke Workshop report. *J Pediatr*
43
44 *2008;153:12e5*.
45
46
47
48 44. Lawrence R, Mathur A, Nguyen The Tich S, et al. A pilot study of continuous limited-
49
50 channel aEEG in term infants with encephalopathy. *J Pediatr* 2009;154:835-841.e1.
51
52
53
54 45. Weiner SP, Painter MJ, Geva D et al. Neonatal Seizures: Electroclinical dissociation.
55
56 *Pediatric Neurol* 1991;7:363-368.
57
58
59
60 46. Hahn CD, Riviello JJ. Neonatal Seizures and EEG. *NeoReviews* 2004;5:e350-e355.

- 1
2
3 47. Mathieson SR, Livingstone V, Low E, et al. Phenobarbital reduces EEG amplitude and
4
5 propagation of neonatal seizures but does not alter performance of automated seizure
6
7 detection. *Clin Neurophysiol* 2016;127:3343-3350.
8
9
10
11 48. Commission of Classification and Terminology of the International League Against
12
13 Epilepsy. Proposal for revised clinical and electroencephalographic classification of
14
15 epileptic seizures. *Epilepsia* 1981;22:489–501.
16
17
18
19 49. Engel J Jr. A Report of the ILAE Classification Core Group. *Epilepsia* 2006;47:1558-1568.
20
21
22
23 50. Nagarajan L, Ghosh S, Palumbo L. Ictal EEG in neonatal seizures: characteristics and
24
25 associations. *Pediatr Neurol* 2011;45:11e6.
26
27
28
29 51. Fisher RS, Cross JH, D'Souza C, et al. Instruction manual for the ILAE 2017 operational
30
31 classification of seizure types. *Epilepsia* 2017;58:531-542.
32
33
34
35 52. Vecchi M, Suppiej A, Mastrangelo M, et al. Focal motor seizure with automatisms in a
36
37 newborn. *Epileptic Dis* 2007;9:149-152.
38
39
40
41 53. Selton D, André M, Hascoët JM. [EEG and ischemic stroke in full-term newborns].
42
43 *Neurophysiol Clin* 2003;33:120-129.
44
45
46
47 54. Low E, Mathieson SR, Stevenson NJ, et al. Early postnatal EEG features of perinatal
48
49 arterial ischaemic stroke with seizures. *PLoS One* 2014;9:e100973.
50
51
52
53 55. Cusmai R, Martinelli D, Moavero R, et al. Ketogenic diet in early myoclonic
54
55 encephalopathy due to non ketotic hyperglycinemia. *Eur J Pediatr Neurol* 2012;16:509 -
56
57 513.
58
59
60

- 1
2
3 56. Ohtahara S, Yamatogi Y. Ohtahara syndrome: with special reference to its
4
5 developmental aspects for differentiating from early myoclonic encephalopathy.
6
7
8 Epilepsy Res 2006;70 Suppl 1:S58-67.
9
10
11 57. Milh M , Villeneuve N, Chouchane M, et al. Epileptic and nonepileptic features in
12
13 patients with early onset epileptic encephalopathy and STXBP1 mutations. *Epilepsia*
14
15 2011;52:1828–1834.
16
17
18
19 58. Porri S, Fluss J, Plecko B, et al. Positive outcome following early diagnosis and treatment
20
21 of Pyridoxal-5'-Phosphate Oxidase Deficiency: a case report. *Neuropediatrics*
22
23 2014;45:64–68.
24
25
26
27 59. Dalla Bernardina B, Aicardi J, Goutières F, et al. Glycine encephalopathy. *Neuropadiatrie*
28
29 1979;10:209-225.
30
31
32
33 60. Cirillo M, Venkatesan C, Millichap JJ, et al. Case report: intravenous and oral pyridoxine
34
35 trial for diagnosis of pyridoxine-dependent epilepsy. *Pediatrics* 2015;136:e257.
36
37
38
39 61. Guerin A, Aziz AS, Mutch C, et al. Pyridox(am)ine-5-Phosphate Oxidase deficiency
40
41 treatable cause of neonatal epileptic encephalopathy with burst suppression: case
42
43 report and review of the literature. *J Child Neurol* 2015;30:1218-1225.
44
45
46
47 62. Simonetti BG, Rieubland C, Courage C, et al. Duplication of the sodium channel gene
48
49 cluster on 2q24 in children with early onset epilepsy. *Epilepsia* 2012;53:2128–2134.
50
51
52
53 63. Numis AL, Angriman M, Sullivan JE, et al. KCNQ2 encephalopathy: Delineation of the
54
55 electroclinical phenotype and treatment response. *Neurology* 2014;82:368-370.
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
64. Plouin P. Benign idiopathic neonatal convulsions (familial and non-familial). In Roger J, Dravet C, Bureau M, et al (Eds) *Epileptic Syndromes in Infancy, Childhood and Adolescence*. 2nd Ed. London: John Libbey, 1992:3-11.
65. Pisano T, Numis AL, Heavin SB, et al. Early and effective treatment of KCNQ2 encephalopathy. *Epilepsia* 2015;56:685–691.
66. Sirsi D, Nadiminti L, Packard MA, et al. Apneic seizures: a sign of temporal lobe hemorrhage in full-term neonates. *Pediatr Neurol* 2007;37:366-370.
67. Scher MS, Aso K, Beggarly ME, et al. Electrographic seizures in preterm and full-term neonates: clinical correlates, associated brain lesions, and risk for neurologic sequelae. *Pediatrics* 1993;91:128-134.
68. Ronen GM, Rosales TO, Connolly M, et al. Seizure characteristics in chromosome 20 benign familial neonatal convulsions. *Neurology* 1993;43:1355-1360.
69. Weckhuysen S, Mandelstam S, Suls A, et al. KCNQ2 encephalopathy: emerging phenotype of a neonatal epileptic encephalopathy. *Ann Neurol* 2012;71:15-25.
70. Commission on Classification and Terminology of the International League Against Epilepsy. Proposal for revised classification of epilepsies and epileptic syndromes. *Epilepsia* 1989;30:389–399.
71. Cornet MC, Sands TT, Cilio MR. Neonatal epilepsies: Clinical management. *Semin Fetal Neonatal Med* Epub 2018 Jan 31.

- 1
2
3 72. Djukic A, Lado FA, Shinnar S, et al. Are early myoclonic encephalopathy (EME) and the
4
5 Ohtahara syndrome (EIEE) independent of each other? *Epilepsy Res* 2006;70 Suppl
6
7 1:S68-76.
8
9
10
11 73. Germano IM, Sperber EF, Ahuja S, et al. Evidence of enhanced kindling and hippocampal
12
13 neuronal injury in immature rats with neuronal migration disorders. *Epilepsia*
14
15 1998;39:1253-1260.
16
17
18
19 74. Raimondi F, Mills P, Clayton PT, et al. A preterm neonate with seizures unresponsive to
20
21 conventional treatment. *BMJ Case Reports Epub* 2015 May 14.
22
23
24
25 75. Schultze S, Weber P, Luetschg J, et al. Incidence and diagnosis of unilateral arterial
26
27 cerebral infarction in newborn infants. *J Perinat Med* 2005;33:170–175.
28
29
30
31 76. Nunes ML, Martins MP, Barea BM, et al. Neurological outcome of newborns with
32
33 neonatal seizures: a cohort study in a tertiary university hospital. *Arq Neuropsiquiatr*
34
35 2008;66:168-174.
36
37
38
39 77. Machado V, Pimentel S, Pinto F, et al. Perinatal ischemic stroke: a five-year retrospective
40
41 study in a level-III maternity. *Einstein (Sao Paulo)* 2015;13:65-71.
42
43
44
45 78. Watanabe K, Miura K, Natsume J, et al. Epilepsies of neonatal onset: seizure type and
46
47 evolution. *Dev Med Child Neurol* 1999;41:318-322.
48
49
50
51 79. Saitsu H, Yamashita S, Tanaka Y, et al. Compound heterozygous BRAT1 mutations cause
52
53 familial Ohtahara syndrome with hypertonia and microcephaly. *J Human Genet*
54
55 2014;59,687–690.
56
57
58
59
60

- 1
2
3 80. Yamamoto H, Okumura A, Fukuda M. Epilepsies and epileptic syndromes starting in the
4 neonatal period. *Brain Dev* 2011;33:213-220.
5
6
7
8
9 81. Kato M, Yamagata T, Kubota M, et al. Clinical spectrum of early onset epileptic
10 encephalopathies caused by KCNQ2 mutation. *Epilepsia* 2013;54:1282-1287.
11
12
13
14 82. Tramonte JJ, Goodkin HP. Temporal lobe hemorrhage in the full-term neonate
15 presenting as apneic seizures. *J Perinatol* 2004;24:726-729.
16
17
18
19
20 83. Okumura A, Hayakawa F, Kato T, et al. Ictal electroencephalographic findings of neonatal
21 seizures in preterm infants. *Brain Dev* 2008;30:261-268.
22
23
24
25
26 84. Castro Conde JR, González-Hernández T, González Barrios D, et al. Neonatal apneic
27 seizure of occipital lobe origin: continuous video-EEG recording. *Pediatrics*
28 2012;129:e1616-1620.
29
30
31
32
33
34 85. Vesoulis ZA, Inder TE, Woodward LJ, et al. Early electrographic seizures, brain injury, and
35 neurodevelopmental risk in the very preterm infant. *Pediatr Res* 2014;75:564-569.
36
37
38
39
40 86. Naim MY, Gaynor JW, Chen J, et al. Subclinical seizures identified by postoperative
41 electroencephalographic monitoring are common after neonatal cardiac surgery. *J*
42 *Thorac Cardiovasc Surg* 2015;150:169-178.
43
44
45
46
47
48 87. Chong DJ, Hirsch LJ. Which EEG patterns warrant treatment in the critically ill? Reviewing
49 the evidence for treatment of periodic epileptiform discharges and related patterns. *J*
50 *Clin Neurophysiol* 2005;22:79-91.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 88. Blume WT, Lüders HO, Mizrahi E, et al. Glossary of descriptive terminology for ictal
4 semiology: report of the ILAE task force on classification and terminology. *Epilepsia*
5 2001;42:1212-1218.
6
7
8
9
10
11 89. Shewmon DA. What is a neonatal seizure? Problems in definition and quantification for
12 investigative and clinical purposes. *J Clin Neurophysiol* 1990;7:315-368.
13
14
15
16
17 90. Yoo JY, Rampal N, Petroff OA, et al. Brief potentially ictal rhythmic discharges in critically
18 ill adults. *JAMA Neurol* 2014;71:454-462.
19
20
21
22
23 91. Yoo JY, Marcuse LV, Fields MC, et al. Brief Potentially Ictal Rhythmic Discharges [B(I)RDs]
24 in Noncritically Ill Adults. *J Clin Neurophysiol* 2017;34:222-229.
25
26
27
28
29 92. Co JP, Elia M, Engel J Jr, et al. Proposal of an algorithm for diagnosis and treatment of
30 neonatal seizures in developing countries. *Epilepsia* 2007;48:1158-1164.
31
32
33
34
35 93. Bourez-Swart MD, van Rooij L, Rizzo C, et al. Detection of subclinical
36 electroencephalographic seizure patterns with multichannel amplitude-integrated EEG
37 in full-term neonates. *Clin Neurophysiol* 2009;120:1916-1922.
38
39
40
41
42 94. Rennie JM, Chorley G, Boylan GB, et al. Non-expert use of the cerebral function monitor
43 for neonatal seizure detection. *Arch Dis Child Fetal Neonatal Ed* 2004;89:F37-40.
44
45
46
47
48 95. Rakshasbhuvankar A, Paul S, Nagarajan L, et al. Amplitude-integrated EEG for detection
49 of neonatal seizures: a systematic review. *Seizure* 2015;33:90-98.
50
51
52
53
54 96. Claassen J, Mayer SA, Kowalski RG, et al. Detection of electrographic seizures with
55 continuous EEG monitoring in critically ill patients. *Neurology* 2004 25;62:1743-1748.
56
57
58
59
60

1
2
3 97. Abend NS, Gutierrez-Colina AM, Topjian AA, et al. Nonconvulsive seizures are common
4
5 in critically ill children. *Neurology* 2011;76:1071-1077.
6
7

8
9 98. Schreiber JM, Zelleke T, Gaillard WD, et al. Continuous video EEG for patients with acute
10
11 encephalopathy in a pediatric intensive care unit. *Neurocrit Care* 2012;17:31-38.
12
13

14
15 99. Gold JJ, Crawford JR, Glaser C, et al. The role of continuous electroencephalography in
16
17 childhood encephalitis. *Pediatr Neurol* 2014;50:318-323.
18
19

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Figure legends

Figure 1: Common etiologies of neonatal seizures in term infants. Adapted from ^{5,6,8-11}

Figure 2: Proposed diagnostic framework of seizures in the neonatal period including classification of seizures. Adapted from 2017 ILAE seizure classification.³ Neonates present with discrete events suspected to be epileptic seizures or are critically ill (often ventilated, sedated and treated with muscle relaxants in intensive care).

Figure 3: Framework for neonatal seizures. Adapted from 2017 ILAE Framework of the epilepsies.⁴ For the purpose of this paper hypoxic events as separate entity because it is the most common etiology of seizures in this age group. There is no evidence at present that immune processes play a role in seizure etiology in this age group.

* Including perinatal hypoxic-ischemic encephalopathy and other hypoxic events in the neonatal period; ** Including infarction, hemorrhage, brain trauma and brain malformations.

Tables

Type	Description ^{3,51}	Special considerations for neonates
Automatisms	A more or less coordinated motor activity usually occurring when cognition is impaired. This often resembles a voluntary movement and may consist of an inappropriate continuation of preictal motor activity	Typically oral and usually in association with other features. Normal and abnormal behavior in term and preterm infants may mimic ictal automatisms. ^{13,52}
Clonic	Jerking, either symmetric or asymmetric, that is regularly repetitive and involves the same muscle groups	Seizure type best recognized clinically. ^{13,14,53,54}
Epileptic spasms	A sudden flexion, extension, or mixed extension–flexion of predominantly proximal and truncal muscles that is usually more sustained than a myoclonic movement but not as sustained as a tonic seizure. Limited forms may occur: Grimacing, head nodding, or subtle eye movements. May occur in clusters.	Rare. May be difficult to differentiate from myoclonic seizures without EMG channel. ⁵⁵⁻⁵⁷
Myoclonic	A sudden, brief (<100 msec) involuntary single or multiple contraction(s) of muscles(s) or muscle groups of variable topography (axial, proximal limb, distal).	Clinically difficult to differentiate from non-epileptic myoclonus. ⁵⁸⁻⁶¹
Sequential seizure	This term is used in the instruction manual for the ILAE 2017 operational classification of seizure types for events with a sequence of signs, symptoms, and EEG changes at different times. ⁵¹	No predominant feature can be determined, instead the seizure presents with a variety of clinical signs. Several features typically occur in a sequence, often with changing lateralization within or between seizures. ⁶²⁻⁶⁴
Tonic	A sustained increase in muscle contraction lasting a few seconds to minutes.	Usually focal, unilateral or bilateral asymmetric. Generalized tonic posturing is often not of epileptic origin. ^{13,56,63,65}
Autonomic	A distinct alteration of autonomic nervous system function involving cardiovascular, pupillary, gastrointestinal, sudomotor, vasomotor, and thermoregulatory functions.	May involve respiration (apnea). Typically seen with other seizure manifestations. EEG confirmation mandatory. ^{13,66-70}
Behavioral arrest	Arrest (pause) of activities, freezing, immobilization, as in behavior arrest seizure.	May be focal and/or followed by apnea, other autonomic manifestations and motor seizures. ⁶⁷
Unclassified seizure type	Due to inadequate information or unusual clinical features with inability to place in other categories.	

Table 1: Integration with the 2017 ILAE Classification of Seizures and Epilepsies

Seizure type	Modifiers
Automatisms	Unilateral Bilateral asymmetric Bilateral symmetric
Clonic seizures	Focal Multifocal Bilateral
Epileptic spasms	Unilateral Bilateral asymmetric Bilateral symmetric
Myoclonic seizures	Focal Multifocal Bilateral asymmetric Bilateral symmetric
Sequential seizure type	Depending on components
Tonic seizures	Focal Bilateral asymmetric Bilateral symmetric

Table 2: Modifiers of motor seizures in the neonatal period

	Term	Preterm
Total	116	20
Gender		
Male	61	10
Female	56	9
GA mean (wks)	39.2	29
PMA at EEG (wks)	40.2	31
Etiology		
HIE	29	5
Structural		
Vascular	22	9
Cortical malformation	6	1
Genetic	25	
Metabolic		
Inborn errors of metabolism	15	
Acute metabolic	3	
Infectious	11	4
Unknown	6	

Table 3: Demographics of 136 neonates included in the data collection.

Diagnosis		Motor							Non-motor		total
		Auto-matisms	Clonic	Epileptic Spasms	Myoclonic	Sequential	Tonic	Autonomic	Behavioral arrest	Electro-graphic	
HIE	term		5			3	2			19	29
	preterm		1				1			3	5
Vascular	term		11			1		2		9	23
	preterm				2		1			7	10
Cortical malformation	term			1	1	3	1		1	1	8
	preterm									1	1
Genetic	term	1	2			14	11				28
	preterm										
IEM	term		1	3	5	4	2			3	18
	preterm										
Acute metabolic	term		1			1				1	3
	preterm										
Infectious	term		1		1	2				8	12
	preterm					1		1		2	4
Unknown	term		2					1		3	6
	preterm										
Total		1	24	4	9	29	18	4	1	57	147

Table 4: Seizure types according to etiology. HIE, infectious etiologies and seizures in premature infants were usually electrographic whereas the most common seizure type in genetic etiologies was sequential. Highlighted in yellow the most common seizures type for a given etiology. HIE: hypoxic ischemic encephalopathy, IEM: inborn errors of metabolism.

	Clinical context of seizure type	Current data review
Automatisms	Described as common seizure type in HIE and preterm infants. ^{13,52,53,74}	Very rare in isolation (<1%).
Clonic	Described as acute symptom of neonatal stroke or cerebral hemorrhage. ^{13,53,54,75-77}	Typical for vascular etiologies in term infants, also commonly seen in HIE .
Myoclonic	Typical seizure type in early myoclonic encephalopathy, but may be seen in other etiologies too, particularly in genetic epilepsy syndromes and preterm infants . Also seen in HIE and inborn errors of metabolism . ^{13,42,55,56,68,69,72,78}	Typical in inborn error of metabolism.
Sequential seizure	Often seen in channelopathies such as BFNE or KCNQ2 encephalopathy, but may be seen with other etiologies. ^{63,64,69,79}	Typical in genetic etiologies.
Spasms	Described in association with inborn error of metabolism or Ohtahara syndrome, here usually in form of tonic spasms (STXBP1, hemimegalencephaly). ^{42,56,58,60,61,78,80}	Rare, mostly in inborn error of metabolism.
Tonic	Typical seizure type in early infantile epileptic encephalopathy (Ohtahara syndrome), but also seen in other epileptic encephalopathies and genetic neonatal epilepsies (eg KCNQ2 and KCNQ3 mutations). May be seen in HIE. ^{13,56,57,63,65,69,72,78,81}	Typical in genetic etiologies.
Autonomic	Described in intraventricular hemorrhage as well as temporal or occipital lobe lesions, typically as apnea and cyanosis. Also described in Ohtahara syndrome. ^{13,64,66,81-84}	Overall rare in isolation (<5%).
Behavioral arrest	Arrest (pause) of activities, freezing, immobilization.	Very rare in isolation (<1%).
Electrographic only seizures	Typical in preterm infants, HIE (particularly in those with basal ganglia/thalamus injury) and neonates undergoing cardiac surgery . ^{6,11-13,17,85,86}	Most common seizure type in preterm infants, HIE and infectious causes.

Table 5: Clinical relevance of seizure types in the neonatal period: review of literature and own data.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: Common etiologies of neonatal seizures in term infants. Adapted from 5,6,8-11

77x42mm (600 x 600 DPI)

view Only




Figure 2: Proposed diagnostic framework of seizures in the neonatal period including classification of seizures. Adapted from 2017 ILAE seizure classification.³ Neonates present with discrete events suspected to be epileptic seizures or are critically ill (often ventilated, sedated and treated with muscle relaxants in intensive care).

90x69mm (600 x 600 DPI)




Figure 3: Framework for neonatal seizures. Adapted from 2017 ILAE Framework of the epilepsies.⁴ For the purpose of this paper hypoxic events as separate entity because it is the most common etiology of seizures in this age group. There is no evidence at present that immune processes play a role in seizure etiology in this age group.

* Including perinatal hypoxic-ischemic encephalopathy and other hypoxic events in the neonatal period; ** Including infarction, haemorrhage, brain trauma and brain malformations.

88x64mm (600 x 600 DPI)