



1º Jornada

Urgencias pediátricas

HPM




1 Convulsiones febriles






1a Línea: Benzodizepinas

Lorazepam



Diazepam



Midazolam



ORIGINAL ARTICLE

Randomized Trial of Three Anticonvulsant Medications for Status Epilepticus

Jaideep Kapur, M.B., B.S., Ph.D., Jordan Elm, Ph.D., James M. Chamberlain, M.D., William Barsan, M.D., James Cloyd, Pharm.D., Daniel Lowenstein, M.D., Shlomo Shinnar, M.D., Ph.D., Robin Conwit, M.D., Caitlyn Meinzer, Ph.D., Hannah Cock, M.D., Nathan Fountain, M.D., Jason T. Connor, Ph.D., and Robert Silbergleit, M.D., for the NETT and PECARN Investigators*

Fenitoína
20 mg/kg

Ácido Valproico
40 mg/kg

Levetiracetam
60mg/kg

Fenitoína
20 mg/kg

Levetiracetam
40mg/kg

2a Línea:

Fenitoína - Fenobarbital - Ácido Valproico - Levetiracetam

Articles

Levetiracetam versus phenytoin for second-line treatment of paediatric convulsive status epilepticus (ECLIPSE): a multicentre, open-label, randomised trial



Mark D Lyttle, Naomi E A Rainford, Carrol Gamble, Shrouk Messahel, Amy Humphreys, Helen Hickey, Kerry Woolfall, Louise Roper, Joanne Noblet, Elizabeth D Lee, Sarah Potter, Paul Tate, Anand Iyer, Vicki Evans, Richard E Appleton, with support of Paediatric Emergency Research in the United Kingdom & Ireland (PERUKI) collaborative*



¡Gracias!

